TOWARDS A METABOLIC ROBOT CONTROL SYSTEM

Jens Ziegler, Peter Dittrich, and Wolfgang Banzhaf

Department of Computer Science
University of Dortmund
44221 Dortmund, Germany
email: ziegler, dittrich, banzhaf@ls11.informatik.uni-dortmund.de
http://ls11-www.informatik.uni-dortmund.de

INTRODUCTION

Bacteria must be able to detect rapid changes in their environment and to adapt their metabolism to external fluctuations. They monitor their surroundings with membrane-bound and intra-cellular sensors. The regulatory mechanisms of bacteria can be seen as a model for the design of robust control systems based on an artificial chemistry. The capability to process information which is needed to keep autonomous agents surviving in unknown environments is discussed.

Biological cells can be seen formally as information processing systems of high complexity [Lengeler, 1996]. They react to a large number of stimuli which are perceived by sensors. These mostly chemical stimuli are converted by *two-component-systems* into signals that are (i) collected, (ii) computed, (iii) integrated and (iv) transmitted through signal transduction pathways. Biochemical changes of proteins, protein-protein interactions and facilitated diffusion along ordered cell structures play a decisive role.

Cellular signal-transmitting systems are hierarchical and often require communication between hundreds of proteins. Membrane-bound two-component systems (see Fig. 1) provide the connection between the *sensory domain* and an attractant or repellant. This domain is connected via helical transmembrane protein structures with the *regulator domain*. Conformational changes in the sensory domain at the external surface induce changes in the relative positioning of the helical membrane-cytosol interface. For instance, binding of a ligand causes a change of the electrostatic density and hence a rotation that increases the distance between charged groups [Stock and Surette, 1996, Lengeler, 1995].

This reaction is transient, i.e. it ends even though the stimulus might continue. Formally, this adaption is achieved through release of a slow feedback reaction which inhibits the signal. Different pathways interlace and so a hierarchical cellular net-



Figure 1. Ligand binding causes the modulation of the electrostatic properties of the helices and therefore the transition between **active** and **inactive** state.

work emerges with a high capacity to store information [Lengeler, 1995, Conrad, 1992]. The computational functions in such biochemical reaction networks have been analysed by [Arkin and Ross, 1994, Bray and Lay, 1994, Bray and Bourret, 1995] and are supposed to function akin to biochemical neuronal networks [Hjelmfelt et al., 1991, Hellingwerf et al., 1995, Okamoto et al., 1995].

Through encapsulation of correlated functions into compartments and the use of superimposed regulating mechanisms (see Fig. 2) the cell reduces the vast flow of information to a communication between a few clearly arranged functional blocks [Darnell et al., 1993, Maynard Smith and Szathmary, 1995].

The signal processing system of a cell is very robust in its dependence upon the observation of central metabolites and, on the other hand, on the hierarchical division into functional blocks. Therefore it can act as a model for the construction of robust, highly parallel and distributed control systems.

CELLS AS INFORMATION PROCESSING SYSTEMS



Figure 2. The metabolic and the regulatory network of a cell.

The metabolism of a cell can be divided into two levels: (i) the metabolic network, where all bio-chemical reactions occur and (ii) the regulatory network which controls the velocity of the reactions and supervises the transduction pathways (see Fig. 2).

All reaction pathways and all regulatory elements together form a kind of highly connected *molecular network*. Each element of this network is a collector of different stimuli (see Fig. 3).



Figure 3. A node in the molecular network of the cell. Stimuli are collected and transmitted depending on the level of adaption to the specific stimulus. This node can be seen as a more general two-component system.

Signals are transmitted depending on the actual state of the adaptor. They are reduced to binary information: by molecules appearing in only two states: *activated* or *inactivated*. Usually these differ only in the absence or presence of a single atom or group at the molecular level. The regulatory network decides through slow or fast feedback or forward reactions which of the connections of the molecular network are activated or inhibited [Lengeler, 1995, Stanier et al., 1986].

One very interesting information processing system of a bacterium controls its movement. Thus, the rest of this section is devoted to a more detailed discussion of the information flow in the chemotaxis system. *Chemotaxis* is the ability of a bacterium to follow concentration gradients by modulation of tumbling frequency. (Fig. 4).



Figure 4. Information flow in the chemotaxis system of *Escherichia Coli*.

In *E. coli* there are six cytoplasmic transduction proteins: *CheA, CheB, CheR, CheW, CheY and CheZ* [Stock and Surette, 1996]. The increase of CheA, for example, indicates the presence of a carbon source (glucose, fructose, etc.). CheA phosphorylates CheY which binds to the switch complex of the flagellar motor and CheB which closes the feedback loop to reset the receptor. The binding of CheY causes clockwise rotation or, to be more specific, it interrupts counterclockwise rotation. Clockwise rotation of the majority of the flagellar motors causes a re-orientation (*tumble*). In this way, the frequency of tumbling is influenced by the concentration of the carbon source.

The cell thus executes a biased random walk, caused by concentration gradients, towards a more favourable environment. It is assumed [Stock and Surette, 1996, Boos, 1977, Alt, 1994] that chemotactic responses are mediated by a *temporal*, rather than by a *spatial*, sensing mechanism, for the following reasons: (i) the length of a bacterium is too small and it moves too quickly to allow for spatial sensing. (ii) Bacteria even respond to sudden changes in concentrations when the concentration is spatially homogeneous. The decision whether to change the movement direction or not depends on the state of some key-metabolites and constitutes an inherent memory of the cell. For a detailed description of bacterial chemotaxis see for example [Bray et al., 1993, Jones and Aizawa, 1991, Lengeler, 1995, Stock and Surette, 1996].

A SPATIAL MODEL OF THE CELL

Information processing in living systems means transduction of substances or molecular interactions. The reception of a stimulus causes the transmission of substances to inner regions of the cell. They arrive either unchanged via diffusion along ordered cell structures or through a cascade of sequential biochemical reactions. Thus, a changing concentration level of X_i can indicate the reception of Y_i. Since chemical reactions take place at very high velocity the delay between encountering a specific substance and a significant change in concentration of the signalling substance is small. Response to an actual stimulus is very fast.

Compartments

Although a procaryotic bacterium appears to lack a cytoskeletton it must not be seen as a "bag full of enzymes" but as a highly ordered system of interacting molecules which are interconnected and connected to the inner membrane.

A model of spatially continuously distributed substances can be based on nonlinear *partial* differential equations. Replacing the spatial derivatives with finite differences divides the system into subsystems (see Fig. 5) which can be treated as homogeneous regions and modelled by a system of coupled *ordinary* differential equations [Jetschke, 1989]. These subsystems are called *compartments*. The state of a compartment is represented by a concentration vector. A realistic model of a complex biochemical system requires 10^2 - 10^5 compartments. In order to keep the simulation of the system tractable the number of compartments should be reduced. The first experiments with a real robot are performed with only one compartment. A model system of this kind can be called a (formal, algorithmic) *reactor*.

Metadynamics

Describing the dynamics of the reactor leads to a system of coupled differential



Figure 5. Spatial model of the cell. Homogeneous subsystems (*compartments*) are connected through diffusion flux. For further details see text.

equations. In order to avoid the number of reactions becoming infinite there must be a focusing on the most relevant ones. A real reactor only contains a finite number of substances and so a possible reaction can only take place when at least one single molecule is inside the reactor. Simulating the dynamics with differential equations, however, assumes a continuous model even with infinitesimal concentrations of substances.

Since a reactor in reality is not continuous but discrete there must be a concentration threshold representing the presence of a single molecule above which the substance is treated as a reactant. If its concentration be lower, it should not participate in the reactor's dynamics. Thus the system has two different time scales: a fast time scale of the concentration changes and a slower time scale for changing the number of participating substances. This change of the ODE system over time is called *metadynamics* [Bagley and Farmer, 1992]. The metadynamics is additionally influenced by diffusion fluxes from other compartments.

ARTIFICIAL REACTION SYSTEMS

Artificial Chemistry

In a real chemistry the kinetic parameters and efficiencies depend on thermodynamics, quantum mechanics and chemical composition of the molecules. For that reason it is impossible to calculate them exactly in practice. To circumvent the troublesome computation of approximations one can formulate an *artificial chemistry* (AC) [Bagley and Farmer, 1992, Fontana, 1992, Fontana, 1994]. An AC is not able to reproduce all properties of real chemistry but it can produce complex behaviour which can be studied instead. By changing the rules different chemistries can be instantiated and, adding layers of more realistic behaviours, it is possible to reproduce properties of the "real" chemistry.

A straightforward way for building a metabolic controller would be, first, to define an arbitrary AC and then use this chemistry to create a controller composed of certain substances. In the following sections we will define two artificial chemistries to show how the abilities of a metabolic controller are depending on its underlying AC. A more radical approach would be to set up a system with the metabolism emerging from the reactions [Banzhaf, 1994].

Example A: Artificial Polymer Chemistry

We begin our discussion with an artificial *polymer chemistry* which is an AC where only polymerisations reactions exists [Bagley and Farmer, 1992]. A substance is represented by a string $s = (s^1, s^2, \ldots s^n) \in A^n$ composed of characters from the alphabet A. The reaction mechanism is simply defined as the concatenation:

$$(s_i^1, s_i^2, \ldots s_i^{n_i}) + (s_j^1, s_j^2, \ldots s_j^{n_j}) \overset{s_k}{\rightleftharpoons} (s_i^1, s_i^2, \ldots s_i^{n_i}, s_j^1, s_j^2, \ldots s_j^{n_j}) \tag{1}$$

The reaction may be catalyzed by a string s_k with catalytic efficiency ν. The dynamic behaviour of the polymer chemistry should be demonstrated by the example shown in figure 6 with the alphabet $A = A, B$. The catalytic constants k_i' are set to $\nu = 10$. Polymerisation is ten times faster than de-polymerisation ($k_i = 1.0$, $k_{-i} = 0.1$).



Figure 6. Example A: Reaction network for the polymer chemistry. The monomers A and B (food set) will be used as inputs. Dotted arrows indicate catalytic activity.

Details of the reaction mechanism and its system of ordinary differential equations (ODE) can be found in the Appendix, part A. The continuous influx flow $\Phi(A)$ and $\Phi(B)$ of the monomers A and B is defined as input. Fig. 7 shows the steady-state input-output relation of the example reaction network.

Example B: Artificial Enyzme-Substrate-Chemistry

The reaction network in Fig. 8 is an example based on enzyme-substrate kinetics. Reaction systems of this kind are referred to as chemical neuronal networks [Hjelmfelt et al., 1991, Okamoto et al., 1995]. The input is represented by the substances C and D, the output by T. The substances marked by "*" are held at a constant concentration level. The detailed reaction mechanism and its corresponding ODE model can be found in the Appendix, part B. Fig. 9 shows the steady-state concentrations of the output species T and several internal substances as a function of input C and D.

Comparison

The polymer chemistry shows a very smooth input-output surface in contrast to the sharp surface of the enzyme-substrate example. The smooth surface insinuates that



Figure 7. Example A: Steady-state input-output relation for the polymer chemistry (Fig. 6).



Figure 8. Example B: Reaction network based on enzyme-substrate kinetics. C and D are input species. T is the output. Substances marked by "*" are held constant.

input information is not processed. The state space which can be reached for different inputs is very large. The surface in state space can not be intuitively divided into different regions with different meanings. In the enzyme-substrate example the quasi-continous input information is condensed by mapping it on two regions in state space which can be clearly separated (Fig. 9). Given some concentrations of possible output-substances the input vector can be much more easily reconstructed than in example B. The robustness of the *classification* can be demonstrated by analysing the information passed from reaction network to reaction network. If these networks are considered to be identical the development of a small error in the first classification is shown in Fig. 10. Transfer functions like a) will reduce the classification to a simple *all-or-nothing*. A correct classification is achieved, even with a disturbed input signal. The course of concentrations in Fig. 8 is similar to a) and thus small fluctuations will be reduced. If the transfer function is of type b) or c) the error will be amplified and the



Figure 9. Example B: Steady-state input-output relation of the enzyme-substrate chemistry example (Fig. 8) which shows the function of an logic AND gate.



Figure 10. Amplification and reduction of a small error during repeated iteration.

original classification can not be restored. Some substances in Fig. 7 are of type b) or c) and thus the classification on incoming information is not robust. Additionally, the total information conservation of the polymer chemistry may lead to information destabilisation when the information should be stored in reaction loops, processed or transmitted. Small fluctuations will be amplified resulting in a quick loss of information.

THE ROBOT CONTROLLER

We shall now discuss the application of the above model to robot control. In order to make use of the computational capabilities of a metabolism it will be necessary to connect it to the environment. If it were simulated without any additional fluxes (e.g. sensor information) the system would run through a transient phase until it finally reached a steady-state.

If we connect the model of our reactor with the infrared sensors of a robot and the motor control with the concentration level of some metabolites we shall be able to use the emerging computational capabilities of the metabolism to control the robot's behaviour. The metabolism is driven away from equilibrium through the sensor information flux, so that all concentrations are influenced by the surroundings of the robot. The sensors are connected with different parts of the metabolism. A change of one sensor value may increase or decrease the concentration of a certain species. The resulting dynamics causes a representation of the environment inside the robot's metabolism.



Figure 11. The robot's metabolism based on enzyme-substrate kinetics. L_i are the sensor substances emitted by the ambient light sensors. D_i are emitted by the distance sensors. Wheel motors are controlled by the substances ML and MR.

In Fig. 11 a schematic metabolism is shown based on an enzyme-substrate kinetics which has been used to control a real robot. The robot has 8 proximity and 8 ambient light sensors. Two basic behaviours are integrated into the metabolism: *obstacle avoidance* and *light seeking*. The part of the reaction network mainly concerned with the obstacle avoidance is processing the substances emitted by the proximity sensors. The

light seeking behaviour gets its input from the ambient light sensors.

Experiments (Fig. 12) show, that the metabolism is able to control the robot in the desired way. It is also able to perform behaviour selection in critical situations where obstacle avoidance has to suppress light seeking. Furthermore, the controller is shown to be robust against disturbance of the reactor, which has been simulated by inserting small random amounts of substances. Figure 12 shows an example of an experiment in an artificial environment.



Figure 12. Example run of the robot. A typical trajectory of the robot's movement is shown in the environment. The robot tries to stay in the lighted area while simultaneously avoiding obstacles.

CONCLUDING REMARKS

The behavior of the robot emerges through chemical reactions between metabolites. It is possible for an autonomous robot to get information about its environment by sensors emitting substances into the robot's metabolism. The control system is parallel and distributed and the ability of navigation emerges only through computation with catalytic reactions [Banzhaf et al., 1996].

The increase and decrease of key-metabolites allows to achieve primitive goals such as obstacle avoidance or luminance gradient following (*phototaxis*). By adding different sensors it would be possible to solve more complicated tasks by stimulus-specific chemical reactions and alarmones. The current focus in our work lies in developing an algorithm which automatically evolves the metabolism to get a better stimulus-response mechanism, but this work is still in progress.

ACKNOWLEDGMENT

This project is supported by the DFG (Deutsche Forschungsgemeinschaft), grant Ba 1042/2-1.

APPENDIX

A: ODE System for Polymer Chemistry

The reaction mechanism for Example A:

$$
\begin{array}{rcll}
A + B & \rightleftharpoons & AB, & \omega_1 = k_1[A][B] - k_{-1}[AB] \\
AB + AB & \rightleftharpoons & ABAB, & \omega_2 = k_2[AB][AB] - k_{-2}[ABAB] \\
ABAB + B & \rightleftharpoons & ABABB, & \omega_3 = k_3[ABAB][B] - k_{-3}[ABABB] \\
A + ABABB & \rightleftharpoons & AABABB, & \omega_4 = k_4[A][ABABB] - k_{-4}[AABABB] \\
A + B & \overset{B}{\rightleftharpoons} & AB, & \omega_5 = k_5\,\nu[A][B][B] - k_{-5}\,\nu[AB][B] \\
AB + AB & \overset{ABAB}{\rightleftharpoons} & ABAB, & \omega_6 = k_6\,\nu\,[AB][AB][ABAB] - k_{-6}\,\nu\,[ABAB][ABAB] \\
ABAB + B & \overset{A}{\rightleftharpoons} & ABABB, & \omega_7 = k_7\,\nu[ABAB][B][A] - k_{-7}\,\nu[ABABB][A] \\
A + ABABB & \overset{AB}{\rightleftharpoons} & AABABB, & \omega_8 = k_8\,\nu\,[A][ABABB][AB] - k_{-8}\,\nu[AABABB][AB]
\end{array}
$$

Table 1. Kinetic rate constants of the polymer chemistry.

parameter	value
k_i	1.0 $\forall i$
k_{-i}	0.1 $\forall i$
ν	10

Resulting ODE system:

$$
\begin{aligned}
\frac{[A]}{dt} &= -\omega_1 - \omega_4 - \omega_5 - \omega_8 + \varphi_A - [A]\,\frac{\varphi}{C} \\
\frac{[B]}{dt} &= -\omega_1 - \omega_3 - \omega_5 - \omega_7 + \varphi_B - [B]\,\frac{\varphi}{C} \\
\frac{[AB]}{dt} &= \omega_1 - \omega_2 + \omega_5 - \omega_6 - [AB]\,\frac{\varphi}{C} \\
\frac{[ABAB]}{dt} &= \omega_2 - \omega_3 + \omega_6 - \omega_7 - [ABAB]\,\frac{\varphi}{C} \\
\frac{[ABABB]}{dt} &= \omega_3 - \omega_4 + \omega_7 - \omega_8 - [ABABB]\,\frac{\varphi}{C} \\
\frac{[AABABB]}{dt} &= \omega_4 + \omega_8 - [AABABB]\,\frac{\varphi}{C}, \qquad (2)
\end{aligned}
$$

with the size C of the reactor. The influx φ is defined as $\varphi = \Phi(A) + \Phi(B)$.

B: ODE System for Enzyme-Substrate Chemistry

The reaction mechanism for Example B:

$$\begin{aligned}
I1^* + C &\rightleftharpoons X1 + C, & \omega_1 &= k_1[C][I1^*] - k_{-1}[C][X] \\
X1 + B &\rightleftharpoons X2^* + A, & \omega_2 &= k_2[X1][B] - k_{-2}[A][X2^+] \\
X3 + A &\rightleftharpoons X4^* + B, & \omega_3 &= k_3[X3][A] - k_{-3}[B][X4^*] \\
X3 &\rightleftharpoons I2^*, & \omega_4 &= k_4[X3] - k_{-4}[I2^*]
\end{aligned}$$

Table 2. Kinetic rate constants of the enzyme-substrate mechanism.

parameter	value	parameter	value
k_1, k_5	70	k_{-1}, k_{-5}	1
k_2, k_6	$5 \cdot 10^4$	k_{-2}, k_{-6}	1
k_3, k_7	$5 \cdot 10^4$	k_{-3}, k_{-7}	1
k_4, k_8	1	k_{-4}, k_{-8}	100

Resulting ODE system:

$$\begin{aligned}
\frac{d[X1]}{dt} &= \omega_1 - \omega_2 \\
\frac{d[X3]}{dt} &= \omega_3 - \omega_4 \\
\frac{d[A]}{dt} = -\frac{[B]}{dt} &= \omega_2 - \omega_3,
\end{aligned} \tag{3}$$

REFERENCES

[Alt, 1994] Alt, W. (1994). Cell motion and orientation. In *Frontiers in mathematical Biology*, pages 79 – 101. Springer.

[Arkin and Ross, 1994] Arkin, A. and Ross, J. (1994). Computational fuctions in biochemical reaction networks. *Biophysical Journal*, 67:560–578.

[Bagley and Farmer, 1992] Bagley, R. J. and Farmer, J. D. (1992). Spontaneous emergence of a metabolism. In Langton, C. G., editor, *Artificial Life II, Proceedings*, Cambridge, MA. MIT Press.

[Banzhaf, 1994] Banzhaf, W. (1994). Self-organization in a system of binary strings. In Brooks, R. A. and Maes, P., editors, *Artificial Life IV, Proceedings*, pages 108 – 119, Cambridge, MA. MIT Press.

[Banzhaf et al., 1996] Banzhaf, W., Dittrich, P., and Rauhe, H. (1996). Emergent computation by catalytic reactions. *Nanotechnology*, 7:307 – 314.

[Boos, 1977] Boos, W. (1977). *Intelligente Bakterien: Chemotaxis als primitives Mittel von Reizleitungssystemen*. Universitätsverlag Konstanz.

[Bray and Bourret, 1995] Bray, D. and Bourret, R. B. (1995). Computer analysis of the binding reactions leading to a transmembrane receptor-linked multiprotein complex involved in bacterial chemotaxis. *Molecular Biology of the Cell*, 6:1367 – 1380.

[Bray et al., 1993] Bray, D., Bourret, R. B., and Simon, M. I. (1993). Computer simulation of the phosphorylation cascade controlling bacterial chemotaxis. *Molecular Biology of the Cell*, 4:469 – 482.

[Bray and Lay, 1994] Bray, D. and Lay, S. (1994). Computer simulated evolution of a network of cell-signalling molecules. *Biophysical Journal*, 66:972 – 977.

[Conrad, 1992] Conrad, M. (1992). Molecular computing: The key-lock paradigm. *Computer*, 25:11–22.

[Darnell et al., 1993] Darnell, J., Lodish, H., and Baltimore, D. (1993). *Molekulare Zellbiologie*. de Gruyter.

[Fontana, 1992] Fontana, W. (1992). Algorithmic chemistry. In Langton, C. G., editor, *Artificial Life II, Proceedings*, Cambridge, MA. MIT Press.

[Fontana, 1994] Fontana, W. (1994). *Molekulare Semantik - Evolution zwischen Variation und Konstruktion*. Rowohlt Verlag.

[Hellingwerf et al., 1995] Hellingwerf, K. J., Postma, P. W., Tommassen, J., and Westerhoff, H. W. (1995). Signal transduction in bacteria: phospho-neural network(s) in escherichia coli? *Federation of Microbiological Societies. Microbiology Review*, 16:309 – 321.

[Hjelmfelt et al., 1991] Hjelmfelt, A., Weinberger, E. D., and Ross, J. (1991). Chemical implementation of neural networks and turing machines. *Proc. Natl. Acad. Sci. USA*, 88:10983–10987.

[Jetschke, 1989] Jetschke, G. (1989). *Mathematik der Selbstorganisation*. Vieweg Verlag.

[Jones and Aizawa, 1991] Jones, C. and Aizawa, S. (1991). The bacterial flagellum and flagellar motor: Structure, assembly and function. *Advances in microbial physiology*, 32:108 –171.

[Lengeler, 1995] Lengeler, J. W. (1995). Basic concepts of microbial physiology and metabolic control. Manuscipt.

[Lengeler, 1996] Lengeler, J. W. (1996). Die biologische Zelle als komplexes Informationssystem. *Biologie in unserer Zeit*, 426:65 – 70.

[Maynard Smith and Szathmary, 1995] Maynard Smith, J. and Szathmary, E. (1995). *The Major Transitions in Evolution*. W. H. Freeman, Oxford.

[Okamoto et al., 1995] Okamoto, M., Tanaka, K., Maki, Y., and Yoshida, S. (1995). Information processing of neural network system composed of 'biochemical neuron': Recognition of pattern similarity in time-variant external analog signals. In Paton, R., editor, *Information Processing in Cells and Tissues. Proceedings.*

[Stanier et al., 1986] Stanier, R. Y., Adelberg, E. A., and Ingraham, J. L. (1986). *The Microbial World*. Prentice-Hall.

[Stock and Surette, 1996] Stock, J. B. and Surette, M. G. (1996). Chemotaxis. In *Escherichia coli and Salmonella typhimurium*, pages 1103 – 1129. American Society for Microbiology, ASM Press.