


COMP 1002

Logic for Computer Scientists

Lecture 18





Cartesian products

- **Cartesian product** of A and B is a set of all pairs of elements with the first from A, and the second from B:



- $A \times B = \{(x, y) | x \in A, y \in B\}$

- $A = \{1, 2, 3\}, B = \{a, b\}$

- $A \times B = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$

- $A = \{1, 2\}, A \times A = \{(1, 1), (1, 2), (2, 1), (2, 2)\}$

| | a | b |
|---|-------|-------|
| 1 | (1,a) | (1,b) |
| 2 | (2,a) | (2,b) |
| 3 | (3,a) | (3,b) |



- Order of pairs does not matter, order within pairs does:
 $A \times B \neq B \times A$.


- Number of elements in $A \times B$ is $|A \times B| = |A| \cdot |B|$

- Can define the Cartesian product for any number of sets:

- $A_1 \times A_2 \times \dots \times A_k = \{(x_1, x_2, \dots, x_k) | x_1 \in A_1 \dots x_k \in A_k\}$

- $A = \{1, 2, 3\}, B = \{a, b\}, C = \{3, 4\}$


- $A \times B \times C = \{(1, a, 3), (1, a, 4), (1, b, 3), (1, b, 4), (2, a, 3), (2, a, 4), (2, b, 3), (2, b, 4), (3, a, 3), (3, a, 4), (3, b, 3), (3, b, 4)\}$





Relations

- **A relation** is a subset of a Cartesian product of sets.
 - If of two sets (set of pairs), call it a **binary** relation.
 - Of 3 sets (set of triples), **ternary**. Of k sets (set of tuples), **k-ary**
- $A=\{1,2,3\}$, $B=\{a,b\}$
 - $A \times B = \{(1, a), (1, b), (2, a), (2, b), (3, a), (3, b)\}$
 - $R = \{(1,a), (2,b),(3,a), (3,b)\}$ is a relation. So is $R=\{(1,b)\}$.
- $A=\{1,2\}$,
 - $A \times A = \{(1,1), (1,2), (2,1), (2,2)\}$
 - $R=\{(1,1), (2,2)\}$ (all pairs (x,y) where $x=y$)
 - $R=\{(1,1),(1,2),(2,2)\}$ (all pairs (x,y) where $x \leq y$).
- $A=PEOPLE$
 - $COUPLES = \{(x,y) \mid \text{Loves}(x,y)\}$
 - $PARENTS = \{(x,y) \mid \text{Parent}(x,y)\}$
- $A=PEOPLE$, $B=DOGS$, $C=PLACES$
 - $WALKS = \{(x,y,z) \mid x \text{ walks } y \text{ in } z\}$
 - Jane walks Buddy in Bannerman park.



Graph of R (bipartite)




Graph of $\{(1,1),(2,2)\}$





Databases and predicates



| Relation R | Predicate P |
|---------------------------------------------------------------------|-----------------------------------------------------|
| A set of tuples | True/false on a given tuple |
| $R = \{(x_1, \dots, x_k) \mid P(x_1, \dots, x_k) \text{ is true}\}$ | $P(x_1, \dots, x_k) \equiv (x_1, \dots, x_k) \in R$ |

- In a database, store relations as tables.
- Then ask queries as predicate logic formulas
 - Return the set of all database elements satisfying the formula.

ProfData

| | A | B | C |
|---|----------|--------------|---------|
| 1 | Manrique | Mata-Montero | EN-2033 |
| 2 | Sharene | Bungay | ER-6032 |
| 3 | Antonina | Kolokolova | ER-6033 |

CourseData

| | A | B | C | D | E |
|---|----------|-----|-------------|---------|--------------|
| 1 | COMP1000 | MWF | 11:00-11:50 | EN-1054 | Mata-Montero |
| 2 | COMP1001 | MWF | 12:00-12:50 | EN-2040 | Bungay |
| 3 | COMP1002 | MTR | 13:00-13:50 | EN-2007 | Kolokolova |

- “Return first names of all profs who teach MWF “
- $Q(fn)$:


$$\exists ln \exists o \text{ ProfData}(fn, ln, o)$$

$$\wedge \exists c, t, r \text{ CourseData}(c, \text{"MWF"}, t, r, ln)$$




Functions

- **A function** $f: X \rightarrow Y$ is a relation on $X \times Y$ such that for every $x \in X$ there is at most one $y \in Y$ for which (x, y) is in the relation.
 - Usual notation: $f(x) = y$
 - y is an **image** of x under f .
 - X is the **domain** of f
 - Y is the **codomain** of f
 - **Range** of f (**image** of X under f):
 - $\{y \in Y \mid \exists x \in X, f(x) = y\}$
 - **Preimage** of a given $y \in Y$:
 - $\{x \in X \mid f(x) = y\}$
 - Preimage of b is $\{2, 3\}$.



This R is not a function




This R is a function with domain $\{1, 2, 3, 4\}$, codomain $\{a, b, c\}$ and range $\{a, b\}$




Functions

- **A function $f: X \rightarrow Y$ is**
 - **Total:** $\forall x \in X \exists y \in Y f(x) = y$
 - $f: \mathbb{Z} \rightarrow \mathbb{Z}$
 - $f(x) = x + 1$ is total.
 - $f(x) = \frac{100}{x}$ is not total.
 - **Onto:** $\forall y \in Y \exists x \in X f(x) = y$
 - $f(x) = x + 1$ is onto over \mathbb{Z} , but not over \mathbb{N}
 - $f(x) = 5x$ is not onto (\mathbb{Z})
 - **One-to-one:** $\forall x_1, x_2 \in X f(x_1) = f(x_2) \rightarrow x_1 = x_2$
 - $f(x) = x + 1$ is one-to-one.
 - $f(x) = x^2$ is not one-to-one
 - **Bijection:** both one-to-one and onto.
 - $f(x) = x + 1$ is a bijection over \mathbb{Z} .


Not total




Not onto



Not one-to-one





Bijection





Functions

- An **inverse** of f is $f^{-1}: Y \rightarrow X$, such that $f^{-1}(y) = x$ iff $f(x) = y$
 - $f(x) = x + 1, f^{-1}(y) = y - 1$
 - Only one-to-one functions have an inverse
- **Composition** of $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ is $g \circ f: X \rightarrow Z$ such that $(g \circ f)(x) = g(f(x))$
 - $f(x) = \frac{x}{5}, g(x) = \lceil x \rceil$, over \mathbb{R}
 - $\lceil x \rceil$ is ceiling: x rounded up to nearest integer.




$$- (g \circ f)(x) = g(f(x)) = \left\lceil \frac{x}{5} \right\rceil$$

$$- (f \circ g)(x) = f(g(x)) = \frac{\lceil x \rceil}{5}$$

$$- (g \circ f)(12.5) = \lceil 2.5 \rceil = 3. \quad (f \circ g)(12.5) = \frac{13}{5} = 2.6$$

- Order matters!





Puzzle: the barber club

- In a certain barber's club,
 - Every member has shaved at least one other member
 - No member shaved himself
 - No member has been shaved by more than one member
 - There is a member who has never been shaved.
- *Question: how many barbers are in this club?*

