



COMP 1002

Logic for Computer Scientists

Lecture 21



Admin stuff

- Assignment 3 is posted
 - Due Monday, March 13
- For the lab this Wednesday, please read definitions in slides for Lecture 19
 - which we did not have time to cover on Tuesday before the midterm





Sum of numbers formula

- Claim: for any $n \in \mathbb{N}$, $\sum_{i=0}^n i = \frac{n(n+1)}{2}$

- Proof.

- Suppose not.

- Let S be a set of all numbers n' such that $\sum_{i=0}^{n'} i \neq \frac{n'(n'+1)}{2}$. By well-ordering principle, if $S \neq \emptyset$, then there is the least number k in S .

- Case 1: $k=0$. But $\sum_{i=0}^0 i = 0 = \frac{0(0+1)}{2}$. So formula works for $k=0$.

- Case 2: $k>0$. Then $k - 1 \geq 0$.

- So $\sum_{i=0}^k i = (\sum_{i=0}^{k-1} i) + k$.

- As k is the smallest bad number, the formula works for $k-1$.

- So $\sum_{i=0}^{k-1} i = \frac{(k-1)k}{2}$

- Now, $\sum_{i=0}^k i = (\sum_{i=0}^{k-1} i) + k = \frac{(k-1)k}{2} + k = \frac{k^2 - k + 2k}{2} = \frac{k^2 + k}{2} = \frac{k(k+1)}{2}$

- So the formula works for $k>0$, too.

- Contradiction. So S is empty, thus the formula works for all $n \in \mathbb{N}$.

Gauss' proof:

$$1 + 2 + \dots + 99 + 100 +$$

$$100 + 99 + \dots + 2 + 1 =$$

$$101 + 101 + \dots + 101 + 101 = 100 \cdot 101$$


$$\text{So } 1+2+ \dots + 99 + 100 = \frac{100 \cdot 101}{2}$$

Works for any n , not just $n=100$



Mathematical induction

- Want to prove a statement $\forall x \in \mathbb{N} P(x)$.
 - Check that $P(0)$ holds
 - And whenever $P(k)$ does not hold for some k , $P(k - 1)$ does not hold either
 - Contradicting well-ordering principle.
 - Contrapositive:
 - if $P(k-1)$ holds for arbitrary k ,
 - then $P(k)$ also must be true.
 - Conclude that $\forall x \in \mathbb{N} P(x)$





Mathematical induction

- Want to prove a statement $\forall x \in \mathbb{N} P(x)$.

- Check that $P(0)$ holds

Proving that $P(0)$ holds is called the **base case**.

- And whenever $P(k)$ does not hold for some k , $P(k - 1)$ does not hold either

- Contradicting well-ordering principle.

- Contrapositive:

That $P(k-1)$ holds is an **induction hypothesis**

- if $P(k-1)$ holds for arbitrary k ,
- then $P(k)$ also must be true.

Proving that $P(k-1) \rightarrow P(k)$ is the **induction step**

- Conclude that $\forall x \in \mathbb{N} P(x)$


Mathematical Induction principle:

If $P(0) \wedge \forall k \in \mathbb{N} P(k) \rightarrow P(k+1)$ then $\forall x \in \mathbb{N} P(x)$



Sum of numbers formula



- Claim: for any $n \in \mathbb{N}$, $\sum_{i=0}^n i = \frac{n(n+1)}{2}$
 - Proof (by induction).
 - $P(n)$ is $\sum_{i=0}^n i = \frac{n(n+1)}{2}$ (*statement* we are proving by induction on n)
 - *Base case*: $k=0$. Then $\sum_{i=0}^0 i = 0 = \frac{0(0+1)}{2}$.
 - *Induction hypothesis*: Assume that $\sum_{i=0}^{k-1} i = \frac{(k-1)k}{2}$ for an arbitrary $k > 0$
 - That is, for an arbitrary number $k-1 \in \mathbb{N}$
 - Can take k instead of $k-1$, but $k-1$ makes calculations simpler.
 - *Induction step*: show that $P(k-1)$ implies $P(k)$.
 - $\sum_{i=0}^k i = (\sum_{i=1}^{k-1} i) + k$.
 - By induction hypothesis, $\sum_{i=1}^{k-1} i = \frac{(k-1)k}{2}$
 - Now, $\sum_{i=1}^k i = (\sum_{i=1}^{k-1} i) + k = \frac{(k-1)k}{2} + k = \frac{k^2 - k + 2k}{2} = \frac{k^2 + k}{2} = \frac{k(k+1)}{2}$
 - *By induction*, therefore, $P(n)$ holds for all $n \in \mathbb{N}$.
- 



Changing the base case

- Mathematical Induction principle:
 - $(P(0) \wedge \forall k \in \mathbb{N} P(k) \rightarrow P(k+1)) \rightarrow \forall x \in \mathbb{N} P(x)$
- What if want to prove it only for $x \geq a$?
 - Make a the base case (when $a \geq 0$). For the rest, assume $k \geq a$.
 - $(P(a) \wedge \forall k \geq a P(k) \rightarrow P(k+1)) \rightarrow \forall x \geq a P(x)$
 - Here, $\forall x \geq a P(x)$ is a shorthand for $\forall x \in \mathbb{N} (x \geq a \rightarrow P(x))$
 - To prove it works, prove $P(n')$ where $n'=n-a$.
- Example: show that for all $n \geq 4$, $2^n \geq n^2$
 - $P(n)$: $2^n \geq n^2$
 - Base case: $n=4$. $2^4 = 16 = 4^2$
 - Induction hypothesis: assume that for an arbitrary $k \geq a$, $2^k \geq k^2$
 - Induction step: show that $2^k \geq k^2$ implies $2^{k+1} \geq (k+1)^2$
 - $2^{k+1} = 2 \cdot 2^k = 2^k + 2^k \geq k^2 + k^2$
 - $(k+1)^2 = k^2 + 2k + 1$.
 - Want: $k^2 + k^2 \geq k^2 + 2k + 1$, so $k^2 \geq 2k + 1$
 - Dividing both sides of the inequality by k : $k \geq 2 + \frac{1}{k}$
 - Since $k \geq 4$, and $2 + \frac{1}{k} \leq 3$, $2 + \frac{1}{k} \leq 3 < 4 \leq k$. So $k \geq 2 + \frac{1}{k}$ and thus $k^2 \geq 2k + 1$
 - So $2^{k+1} = 2 \cdot 2^k = 2^k + 2^k \geq k^2 + k^2 \geq k^2 + 2k + 1 = (k+1)^2$
 - By induction, for all $n \geq 4$, $2^n \geq n^2$
- Corollary: as n grows, an algorithm running in time n^2 will quickly outperform an algorithm running in time 2^n



Strong induction

- For our coins problem, needed not just $P(k-1)$, but $P(k-3)$, and to look at three cases.
- **Mathematical Induction** principle:
 - $(P(0) \wedge \forall k \in \mathbb{N} P(k) \rightarrow P(k+1)) \rightarrow \forall x \in \mathbb{N} P(x)$
- **Strong Induction** principle:
 - $(\exists b \in \mathbb{N} \forall c \in \mathbb{N} (0 \leq c \wedge c \leq b \rightarrow P(c)))$
 $\wedge \forall k > b (\forall i \in \{0, \dots, k-1\} P(i)) \rightarrow P(k)$
 $\rightarrow \forall x \in \mathbb{N} P(x)$
- Strong induction seems stronger...
 - But in fact, mathematical induction, strong induction and well-order principles are equivalent to each other.
 - So choose the most convenient one.



Puzzle: coins



- A not-too-far-away country recently got rid of a penny coin, and now everything needs to be rounded to the nearest multiple of 5 cents...
 - Suppose that instead of just dropping the penny, they would introduce a 3 cent coin.
 - Like British three pence.
 - What is the largest amount that cannot be paid by using only existing coins (5, 10, 25) and a 3c coin?

7c

Any number $n > 7$ can be paid with 3,5,10,25 coins (even just 3 and 5).



Strong induction

- **Strong Induction** principle (general form):
 - $(\exists b \in \mathbb{N} \forall c \in \mathbb{N} (a \leq c \wedge c \leq b \rightarrow P(c))$
 $\wedge \forall k > b (\forall i \in \{a, \dots, k-1\} P(i)) \rightarrow P(k))$
 $\rightarrow \forall x \in \mathbb{N} (x \geq a \rightarrow P(x))$
- **Coins:** $\forall x \in \mathbb{N}$, if $x > 7$ then $\exists y, z \in \mathbb{N}$ such that $x = 3y + 5z$.
 - $P(n)$: $\exists y, z \in \mathbb{N} n = 3y + 5z$. Also, $a=8$.
 - Base cases: $b = 10$, so $c \in \{8, 9, 10\}$
 - $n=8$. $8 = 3 \cdot 1 + 5 \cdot 1$, so $y=1, z=1$.
 - $n=9$. $9=3 \cdot 3$, $y=3, z=0$
 - $n=10$. $10=5 \cdot 2$, $y=0, z=2$.
 - Induction hypothesis: Let k be an arbitrary integer such that $k > 10$. Assume that for all $i \in \mathbb{N}$ such that $8 \leq i < k \exists y_i, z_i \in \mathbb{N} i = 3y_i + 5z_i$
 - Induction step. Show that induction hypothesis implies that $\exists y, z \in \mathbb{N} k = 3y + 5z$
 - Since $k \geq b$, $k - 3 \geq a$. So by induction hypothesis $\exists y_{k-3}, z_{k-3} \in \mathbb{N} k - 3 = 3y_{k-3} + 5z_{k-3}$. Now take $z=z_{k-3}$ and $y = y_{k-3} + 1$. Then $k = 3y + 5z$.
 - By strong induction, get that for all $x > 7$, $\exists y, z \in \mathbb{N}$ such that $x = 3y + 5z$.



Puzzle: all horses are white



- Claim: all horses are white.
- Proof (by induction):
 - $P(n)$: any n horses are white.
 - Base case: $P(0)$ holds vacuously
 - Induction hypothesis: any k horses are white.
 - Induction step: if any k horses are white, then any $k+1$ horses are white.
 - Take an arbitrary set of $k+1$ horses. Take a horse out.
 - The remaining k horses are white by induction hypothesis.
 - Now put that horse back in, and take out another horse.
 - Remaining k horses are again white by induction hypothesis.
 - Therefore, all the $k+1$ horses in that set are white.
 - By induction, all horses are white.

