


COMP 1002

Intro to Logic for Computer Scientists

Lecture 8





False premises



- An argument can still be valid when some of its premises are false.
 - Remember, false implies anything.
- Bertrand Russell: “If $2+2=5$, then I am the pope”
 - Suppose $2+2=5$
 - If $2+2=5$, then $1=2$ (subtract 3 from both sides).
 - So $1=2$ (by modus ponens)
 - Me and the pope are two people.
 - Since $1=2$, me and the pope are one person.
 - Therefore, I am the pope!

Natural deduction vs. Truth tables

- In this puzzle, it was faster to solve it using modus ponens (natural deduction method) than writing a truth table.
- But is it always better?
- The answer is...

Nobody knows!

- It is a very closely related to the question of how fast can one check if something is a tautology.
 - And that's a million dollar question!





The million dollar question




- In English, known as “P vs. NP” problem
 - P stands for “polynomial time computable”.
 - NP is “polynomial time checkable”
 - non-deterministic polynomial-time computable
 - **Question: is everything efficiently checkable also efficiently computable?**
- In Russian, called “perebor” problem.
 - “perebor” translates as “exhaustive search”.
 - Question: is it always possible to avoid looking through nearly all potential solutions to find an answer?
 - **Are there situations when exhaustive search is unavoidable?**




The million dollar question

- **NP-completeness:** enough to answer for the problem of checking satisfiability (SAT)!
- A formula is like a basket of apples.
formula is a tautology
=
All apples in the basket are good.
- Can you check that all apples are good without looking at every single one?
- Can you do it for every possible basket of apples?
 - Smell test?



Automated provers



- How to make an automated prover which checks whether a formula is a tautology?
 - And so can check if an argument is valid, etc.
- Truth tables:
 - easy to program, but proofs are huge.
- Natural deduction:
 - proofs might be smaller than a truth table
 - Are they always? Good question...
 - even if there is a small proof, how can we find one quickly?
 - Nobody knows...



Resolution proofs



- Middle ground: use the **resolution rule**:
 - Basis for many practical provers (SAT solvers).
 - Used in verification, scheduling, etc...

$$\begin{array}{l} C \vee x \\ D \vee \neg x \end{array}$$

$$\begin{array}{l} y \vee \neg z \vee w \\ u \vee \neg w \end{array}$$

$$\begin{array}{l} y \vee w \vee \neg z \\ \neg z \vee \neg w \end{array}$$

$$\frac{}{\therefore C \vee D}$$

$$\frac{}{\therefore y \vee \neg z \vee u}$$

$$\frac{}{\therefore y \vee \neg z}$$

- Ignore order in an OR and remove duplicates.

Natural deduction



- A: this house is next to a lake.
- B: the treasure is in the kitchen
- C: The tree in front is elm
- D: the treasure is under the flagpole.
- E: The tree in the back is oak
- F: The treasure is in the garage

1. If A then not B
2. If C then B
3. A
4. C or D
5. If E then F
6. Not B
7. Not C
8. D

- If house is next to the lake then the treasure is not in the kitchen
- The house is next to the lake
- Therefore, the treasure is not in the kitchen.

Puzzle 8

- Suppose that nobody in our class carries more than 10 pens.
- There are 70 students in our class.
- Prove that there are at least 2 students in our class who carry the same number of pens.
 - In fact, there are at least 7 who do.

