

CS 2742 (Logic in Computer Science) – Winter 2013

Lecture 5

Antonina Kolokolova

January 18, 2013

1.1 Proof techniques

Example 1. Consider the sentence “if n is divisible by 4, then n is divisible by 2” (we will use the notation $n|4$ to mean n is divisible by 4). This is an if-then statement. Its contrapositive is “if $n \nmid 2$ then $n \nmid 4$. That is, if n is an odd number then it is definitely not divisible by 4. So $n|4$ is sufficient for $n|2$ (if n is divisible by 4, it is sufficient for n to be divisible by 2). On the other hand, $n|2$ is necessary for $n|4$.

- 1) *Direct proof*: show that if p is true directly.
- 2) *Proof by contrapositive*: instead of $p \rightarrow q$ prove $\neg q \rightarrow \neg p$.

Lemma 1. *If n^2 is even, then n is even.*

Proof. We will show this by showing that if n is odd, then n^2 is odd. If n is odd, then $n = 2k + 1$ for some k . Then $(2k + 1)^2 = 2(2k^2 + 2k) + 1$, which is an odd number. This proves that if n is odd, then n^2 is odd, thus proving the contrapositive of if n^2 is even then n is even, and so proving the statement “if n^2 is even then n is even” itself. \square

- 3) *Proof by contradiction*: to show that p is true, show that $\neg p \rightarrow F$.
It is easy to show that $(\neg p \rightarrow F)$ is logically equivalent to p : just note that $(\neg\neg p \vee F) \iff (p \vee F) \iff p$ by applying the definition of implication followed by the double negation law followed by the identity law.

Theorem 1. *$\sqrt{2}$ is irrational.*

Proof. Recall that a number is called *rational* if it can be represented as an (irreducible) fraction of two integers. Assume, for the sake of contradiction, that $\sqrt{2}$ is rational: that is, there are integers m and n which do not have any common divisors > 1 such that $\sqrt{2} = m/n$.

- If $\sqrt{2} = m/n$ then $(\sqrt{2})^2 = m^2/n^2$.
- From here, $2n^2 = m^2$, which means that m^2 is even.
- By the lemma above, then m is even, so $m = 2k$ for some k .
- Then $m^2 = 4k^2$. So $2n^2 = 4k^2$, and, dividing by 2, $n^2 = 2k^2$. So n^2 is even.
- Using the lemma again, conclude that n is even.
- So both n and m are even, but we assumed that m and n do not have a non-trivial common divisor. This is a contradiction.

□

4) *Proof by cases:* to show that p is true, prove $(q \rightarrow p) \wedge (\neg q \rightarrow p)$.

Lemma 2. For any natural number n , $\lfloor (n+1)/2 \rfloor = \lceil n/2 \rceil$

Here, $\lfloor k \rfloor$ is a *floor* of a (real) number, defined to be the largest integer smaller than or equal to k . For example, $\lfloor 5/2 \rfloor = 2$, and $\lfloor 4/2 \rfloor = 2$. Similarly, a *ceiling* of a number is the smallest integer larger than or equal to that number. So $\lceil 5/2 \rceil = 3$ and $\lceil 4/2 \rceil = 2$. For an integer, both its floor and its ceiling are equal to that integer itself; for a non-integer, the floor is its rounded-down value and the ceiling is rounded up.

Proof. Case 1: n is even. Then $\lfloor (n+1)/2 \rfloor = n/2 = \lceil n/2 \rceil$.

Case 2: n is odd. Then $\lfloor (n+1)/2 \rfloor = (n+1)/2 = \lceil n/2 \rceil$.

□

Puzzle 4. A from the island of knights and knaves said: “If I am a knight, then I’ll eat my hat!”. Prove that A will eat his hat.